

(19)



Europäisches Patentamt
European Patent Office
Office européen des brevets

(11) Publication number:

0 263 638
A2

(12)

EUROPEAN PATENT APPLICATION

(21) Application number: 87308657.3

(57) Int. Cl.⁴: A61K 7/16, A61K 7/18

(22) Date of filing: 30.09.87

(30) Priority: 02.10.86 GB 8623643

(43) Date of publication of application:
13.04.88 Bulletin 88/15(84) Designated Contracting States:
AT BE CH DE ES FR GB GR IT LI NL SE(71) Applicant: UNILEVER PLC
Unilever House Blackfriars P.O. Box 68
London EC4P 4BQ(GB)

(84) GB

(71) Applicant: UNILEVER NV
Burgemeester s'Jacobplein 1 P.O. Box 760
NL-3000 DK Rotterdam(NL)

(84) BE CH DE ES FR GR IT LI NL SE AT

(72) Inventor: Clarkson, John Richard
68 Borrowdale Road
Bebington Wirral L63 3AR(GB)
Inventor: Duckworth, Ralph Marsland
45 Kelsborrow Way
Kelsall Nr Chester(GB)
Inventor: Murray, Andrew Malcolm
15 Moorings Close
Parkgate South Wirral Cheshire(GB)
Inventor: Price, Timothy John
48 Garth Drive
Chester CH2 2AG(GB)(74) Representative: Doucy, Robert Henry et al
Unilever PLC Patents Division P.O. Box 68
Unilever House
London EC4P 4BQ(GB)

(54) Oral composition.

EP 0 263 638 A2

(57) An oral hygiene product especially for inhibiting caries is the form of toothpaste or mouthwash. The product maintains a low fluoride ion concentration in the mouth for longer periods than conventional products by the rapid precipitation of calcium fluoride either in the mouth or immediately prior to use. The product comprises a composition containing calcium ions and a further composition containing fluoride ions. The ion product of the calcium and fluoride ions of the combined product preferably exceeds $3 \times 10^{-8} \text{ mol}^3 \text{ dm}^{-9}$.

ORAL PRODUCTS

This invention relates to oral products, more particularly oral hygiene products which are effective to combat dental caries.

It is well known to include water-soluble fluorine-containing salts, for example sodium fluoride or sodium monofluorophosphate, in oral products, especially mouthwashes and toothpastes, and that by regular use of such products the incidence of dental caries can be reduced. It is believed that the fluoride ion or monofluorophosphate ion, interacts with the tooth substance to increase its resistance to acid attack and to aid repair of carious enamel. However, the opportunity for this efficacious interaction to occur is short-lived because the oral fluoride level falls off rapidly after use of a mouthwash or toothpaste.

There have been attempts in the past to provide means for maintaining a certain concentration of fluoride ions in the mouth over a longer period. These have included proposals for locating a fluoride ion source in the mouth, for example as part of an orthodontic appliance.

The Applicants have investigated the possibility of depositing particles of a fluoride ion-releasing material in the mouth and have discovered that surprising results are achieved through the use of particles of freshly precipitated calcium fluoride, as is more particularly described hereinafter.

It is already known to treat teeth to provide anticaries protection with compositions containing calcium ions and fluoride ions. Such compositions are described in EP-A-89 136 (Procter & Gamble) and they comprise a calcium ion source, a fluoride ion source and a calcium sequestering agent possessing specific solubility and binding properties to control the precipitation of calcium fluoride. The calcium-sequestering agent present in these compositions has a stability constant sufficient to inhibit uncontrolled and rapid precipitation of calcium fluoride.

In US-A-4 080 440 (DiGuilo et al) there is described a two-part product for example mouthwash or toothpaste, for remineralising demineralised tooth enamel, consisting of a first solution containing calcium ions and a second solution containing phosphate ions and optionally fluoride ions, which solutions when mixed together form a metastable solution having a pH of about 4 or below. The metastable solution is applied to the tooth surface within 5 minutes of its formation. The ions diffuse into the demineralised subsurface where due to a rise in pH calcium phosphate and calcium fluoride are precipitated.

GB-A-1 452 125 (Procter & Gamble) also describes a two-part oral treatment product comprising a first solution containing calcium ions and a second solution containing phosphate ions and fluoride ions, the solutions being applied to the tooth surface sequentially. The respective ions diffuse to the subsurface dental enamel, the ions of the second applied solution coming into contact with those previously deposited and forming a precipitate which is bound to the tooth structure.

There have also been described in the literature various compositions containing both a calcium compound and a fluoride compound and wherein means are provided to inhibit the formation of calcium fluoride; examples are EP-A-40 738 (Richardson-Vicks), GB-A-777 556 (Colgate-Palmolive) and US-A-4 098 435 (Weyn).

GB 1 090 340 (Warner Lambert) discloses a one-part oral composition containing calcium and fluoride ions. Any calcium fluoride in such compositions will not be freshly precipitated.

GB 1 408 922 (Blendax) and US 4 108 980 (Colgate) disclose compositions comprising calcium, phosphate and fluoride ions for remineralising dental enamel. The disclosed compositions contain such high levels of phosphate that preferential precipitation of solid calcium phosphate phases such as octacalcium phosphate and hydroxyapatite would deplete the liquid phase composition of calcium ions rather than allow calcium fluoride precipitation. Further, any calcium fluoride precipitation would occur slowly due to the presence of large amounts of the crystal growth inhibitor, phosphate.

The present invention provides an oral preparation for inhibiting caries, which comprises as a combined preparation a first composition and a second composition for admixing in the mouth or for admixing immediately prior to introduction into the mouth, wherein

the first composition comprises an aqueous solution containing calcium ions,

the second composition comprises an aqueous solution containing fluoride ions,

the first and second compositions being such that when mixed rapid precipitation of calcium fluoride occurs.

For optimum retention of fluoride in the mouth the ratio of calcium ions to fluoride ions should be stoichiometric (i.e. 1:2).

Rapid precipitation of calcium fluoride is induced by using high concentrations of calcium ions and fluoride ions. Thus, the initial ion product (IP) on mixing the two compositions which is given by the equation:

$$5 \quad IP = (Ca^{2+})(F^{-})^2$$

where (Ca^{2+}) = calcium ion concentration, and

(F^{-}) = fluoride ion concentration

should exceed the solubility product of calcium fluoride, $2.85 \times 10^{-11} \text{ mol}^3 \text{ dm}^{-9}$ at 25°C , by at least 1000 times. More preferably, the above ion product should exceed $3 \times 10^{-7} \text{ mol}^3 \text{ dm}^{-9}$.

Rapid precipitation of calcium fluoride is not favoured by low concentrations of calcium ions and fluoride ions. At low concentrations of the aforementioned ions a metastable solution may be formed so that precipitation proceeds slowly for many minutes after the ions are mixed. Rapid precipitation is also inhibited by crystal growth inhibitors such as hydrogen phosphate ions (HPO_4^{2-}).

15 The presence of substances which react with either calcium ions or fluoride ions to produce material having a solubility less than that of calcium fluoride are not advantageous. The presence of substances which reduce the amount of free calcium or fluoride ions before the compositions are mixed are also not advantageous. Examples of such materials are metal ion sequestrants such as EDTA, which react with calcium ions to form complexes, and alumina abrasive particles which bind fluoride.

20 In preferred embodiments of the invention the composition is in the form of a two-part toothpaste or a two-part mouthwash.

Calcium chloride is used in a preferred embodiment of the invention. Preferably a calcium ion concentration of at least 3.95 mM is used, more preferably at least 8mM. Calcium ion levels in excess of 0.3M provide no additional benefit and would generally not be used.

25 Sodium fluoride is used in a preferred embodiment of the invention. A fluoride ion concentration of at least 7.9mM is preferably used, more preferably at least 16mM. Fluoride ions are preferably less than 0.3M.

One or both of the compositions may contain adjuvants. Such adjuvants may include colouring agents, flavours, humectants, abrasives, detergents, preferably nonionic detergents, and the like and other therapeutic agents compatible with calcium or fluoride ions.

30 In use the first and second compositions are admixed in the mouth or immediately prior to introduction into the mouth. A delivery system providing for physical separation of the two compositions and for simultaneous or sequential delivery of the compositions may also be used.

The efficacy of the freshly precipitated calcium fluoride in maintaining fluoride ion levels in the mouth is demonstrated by the following experiments.

35

EXPERIMENT 1

Procedure

40

The procedure employed for each test was as follows:

(i) a 2ml mouthwash was held in the mouth for one minute and then expectorated. Such mouthwashes consisted of two 1 ml solutions, which were either applied separately or mixed before application.

(ii) saliva samples were collected at regular intervals for several hours after mouthwash application.

45 (iii) Saliva samples were buffered to pH5 by adding 10% by weight of TISAB buffer and then fluoride activities were measured using a fluoride ion specific electrode (Orion 94-09).

The following aqueous solutions were used:

a) 0.0526 M sodium chloride

b) 0.0526 M calcium chloride and 0.1053 M sodium fluoride mixed immediately prior to application

50 c) as b) but mixed 1.5 hours before application.

d) a suspension of the equivalent concentration of calcium fluoride to that of (c) diluted from a concentrated stock prepared 3 months before application. (Aged calcium fluoride).

55

Results

Figure 1 shows that salivary fluoride levels resulting from application of mouthwashes containing freshly precipitated calcium fluoride are significantly higher than the corresponding values for aged precipitates. The mouthwash containing the most aged calcium fluoride particles (three months old) gave salivary fluoride values only slightly greater than the 'baseline' values measured after application of sodium chloride solution.

10 EXPERIMENT 2

The following experiment was performed to demonstrate that a mouthwash containing freshly precipitated calcium fluoride is capable of maintaining fluoride in the mouth at a higher level than a sodium fluoride mouthwash of the same fluoride content.

15 Procedure

The experiment was carried out in a similar manner to that described above using the following aqueous solutions;

- i) 0.0526M calcium chloride and 0.1052M sodium fluoride mixed immediately prior to application
- ii) 0.0526M sodium fluoride.

25 Results

Figure 2 shows that salivary fluoride levels resulting from the application of mouthwashes containing freshly precipitated calcium fluoride are significantly higher than the corresponding values for sodium fluoride.

30 The invention is illustrated by reference to the following examples. All percentages are by weight.

35

40

45

50

55

EXAMPLE 1Mouthwash

First composition

%

Sodium fluoride	0.11
Sorbitol syrup (70% solution)	15.0
Ethanol	10.0
Nonionic detergent	0.4
Saccharin	0.04
Flavour	0.15
Water	to 100

Second composition

%

Calcium chloride dihydrate	0.195
Sorbitol syrup (70% solution)	15.0
Ethanol	10.0
Nonionic detergent	0.4
Saccharin	0.04
Flavour	0.15
Water	to 100

Approximately 5 ml of each component are mixed together immediately before use.

EXAMPLE 25 Toothpaste

	First composition	<u>%</u>
10	Sodium fluoride	0.44
	Abrasive silica	14.0
	Thickening silica	8.0
15	Sorbitol syrup (70% solution)	50.0
	Sodium carboxymethylcellulose	0.65
	Nonionic detergent	1.5
20	Flavour	1.0
	Saccharin	0.1
	Titanium dioxide	1.0
25	Water	to 100

adjust pH to 7.0-7.5 with NaOH.

30

	Second composition	<u>%</u>
35	Calcium chloride dihydrate	0.78
	Abrasive silica	14.0
	Thickening silica	8.0
40	Sorbitol syrup (70% solution)	50.0
	SCMC	0.65
	Nonionic detergent	1.5
45	Flavour	1.0
	Saccharin	0.1
	Titanium dioxide	1.0
	Water	to 100

50 adjust pH to 7.0-7.5 with NaOH.

to be mixed in equal proportions in use.

55 Toothpaste-mouthwash combinations are also within the scope of the invention. One such combination can be achieved by applying the first composition, 1 ml of a mouthwash of composition

%

5	Sodium fluoride	0.44
	Sorbitol syrup (70% solution)	15.0
	Ethanol	10.0
10	Nonionic detergent	0.4
	Saccharin	0.04
	Flavour	0.15
15	Water	to 100.0

and then immediately brushing with the second composition of Example 2.

The oral product of the invention may also be incorporated in other oral preparations or formulations such as dual composition lozenges, sweets and chewing gum.

Claims

1. An oral preparation for inhibiting caries, characterised in that the preparation comprises as a combined preparation a first composition and a second composition for admixing in the mouth or for admixing immediately prior to introduction into the mouth, wherein

the first composition contains a source of calcium ions,

the second composition contains a source of fluoride ions,

the first and second compositions being such that when mixed rapid precipitation of calcium fluoride occurs.

2. A preparation as claimed in Claim 1 wherein the ion product (IP) of the calcium ions and fluoride ions as hereinbefore defined exceeds $3 \times 10^{-8} \text{ mol}^3 \text{ dm}^{-9}$ on mixing.

3. A preparation as claimed in Claim 1 or Claim 2 wherein the calcium salt is calcium chloride.

4. A preparation as claimed in any one of Claims 1 to 3, wherein the fluoride salt is sodium fluoride.

5. A preparation as claimed in any one Claims 1 to 4, wherein the compositions are in the form of a two-part mouthwash.

6. A preparation as claimed in any one of Claims 1 to 4, wherein the compositions are in the form of a two-part toothpaste.





Fig.1

MEAN SALIVA F VALUES FOLLOWING IN SITU CaF_2 PRECIPITATION

Fig.2



THIS PAGE BLANK (USPTO)

⑫

EUROPEAN PATENT APPLICATION

⑲ Application number: 87308657.3

⑤① Int. Cl. 4: **A61K 7/16 , A61K 7/18**

⑳ Date of filing: 30.09.87

③① Priority: 02.10.86 GB 8623643

④③ Date of publication of application:
13.04.88 Bulletin 88/15

⑧④ Designated Contracting States:
AT BE CH DE ES FR GB GR IT LI NL SE

⑧⑧ Date of deferred publication of the search report:
26.04.89 Bulletin 89/17

⑦① Applicant: **UNILEVER PLC**
Unilever House Blackfriars P.O. Box 68
London EC4P 4BQ(GB)

⑧④ **GB**

⑦① Applicant: **UNILEVER NV**
Burgemeester s'Jacobplein 1 P.O. Box 760
NL-3000 DK Rotterdam(NL)

⑧④ **BE CH DE ES FR GR IT LI NL SE AT**

⑦② Inventor: **Clarkson, John Richard**
68 Borrowdale Road
Bebington Wirral L63 3AR(GB)
Inventor: **Duckworth, Ralph Marsland**
45 Kelsborrow Way
Kelsall Nr Chester(GB)
Inventor: **Murray, Andrew Malcolm**
15 Moorings Close
Parkgate South Wirral Cheshire(GB)
Inventor: **Price, Timothy John**
48 Garth Drive
Chester CH2 2AG(GB)

⑦④ Representative: **Doucy, Robert Henry et al**
Unilever PLC Patents Division P.O. Box 68
Unilever House
London EC4P 4BQ(GB)

⑤④ **Oral composition.**

EP 0 263 638 A3

⑤⑦ An oral hygiene product especially for inhibiting caries is the form of toothpaste or mouthwash. The product maintains a low fluoride ion concentration in the mouth for longer periods than conventional products by the rapid precipitation of calcium fluoride either in the mouth or immediately prior to use. The product comprises a composition containing calcium ions and a further composition containing fluoride ions. The ion product of the calcium and fluoride ions of the combined product preferably exceeds $3 \times 10^{-8} \text{ mol}^3 \text{ dm}^{-3}$.



European Patent
Office

EUROPEAN SEARCH REPORT

Application Number

EP 87 30 8657

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int. Cl. 4)
X, P	EP-A-0 222 603 (NEIRINCKX, RUDI DOMINIQUE) * column 4, lines 26 - 30; claims * ---	1-6	A 61 K 7/18 A 61 K 7/16
X	WO-A-8 602 265 (ROLLA, GUNNAR) * page 2, paragraphs 2 - 3 * ---	1-6	
X	FR-A-2 361 851 (COLGATE-PALMOLIVE COMPANY) * page 15, lines 27 - 36; claims * -----	1-2,4,6	
A		3,5	
			TECHNICAL FIELDS SEARCHED (Int. Cl. 4)
			A 61 K
The present search report has been drawn up for all claims			
Place of search BERLIN		Date of completion of the search 20-12-1988	Examiner SIATOU E
CATEGORY OF CITED DOCUMENTS X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons & : member of the same patent family, corresponding document			

EPO FORM 1503 03.82 (P0401)

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ BLACK BORDERS
- ☐ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES
- ☒ FADED TEXT OR DRAWING
- ☒ BLURRED OR ILLEGIBLE TEXT OR DRAWING
- ☐ SKEWED/SLANTED IMAGES
- ☐ COLOR OR BLACK AND WHITE PHOTOGRAPHS
- ☐ GRAY SCALE DOCUMENTS
- ☐ LINES OR MARKS ON ORIGINAL DOCUMENT
- ☐ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY
- ☐ OTHER: _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.

THIS PAGE BLANK (USPTO)